

# Medications for AUD Treatment in Adults

### Withdrawal Management

- Benzodiazepine
  - Chlordiazepoxide (Librium)
  - Diazepam (Valium)
  - Lorazepam (Ativan)
  - Phenobarbital (Luminal)
- GABA-ergics
  - Carbamazepine (Tegretol)
  - Gabapentin (Neurontin)
  - Valproic acid (Depakote)

### Relapse Prevention

- On-Label
  - Acamprosate (Campral)
  - Disulfiram (Antibuse)
  - Naltrexone (ReVia & Vivitrol)
- Off-Label
  - Gabapentin (Neurontin)
  - Topiramate (Topamax)
  - Baclofen (Gablofen, Lioresal)



Chlordiazepoxide (Librium)

Route Oral **Formulation** • 5mg, 10mg, 25mg capsules • Symptom-driven or fixed dose-and-Daily Dose taper protocols Half-Life • Intermediate acting (10-30 hours) Cost • 0.18\$-4.33\$ per capsule



Chlordiazepoxide (Librium)

### Common Side Effects

- Confusion
- Drowsiness
- Restlessness
- Irritability
- Tolerance
- Misuse

#### **Precautions**

- Dose adjustment in elderly
- Psychiatric patients
- Hyperactive aggressive pediatric patients

#### Other

 Preferred for patients with liver dysfunction



Diazepam (Valium)

Route Oral **Formulation** • 2mg, 5mg, 10mg tablets • Symptom-driven or fixed dose-and-Daily Dose taper protocols Half-Life • Long acting (20-100 hours) Cost • 0.14\$-4.88\$ per tablet



Diazepam (Valium)

#### Common Side Effects

- Confusion
- Drowsiness
- Restlessness
- Irritability
- Tolerance
- Misuse

#### Precautions

- Pregnancy
- Other substance use disorder
- Depression
- Seizures
- Glaucoma



Lorazepam (Ativan)

Route Oral Formulation • 0.5mg, 1mg, 2mg tablets • Symptom-driven or fixed dose-and-Daily Dose taper protocols Half-Life • Short acting (10-20 hours) Cost • 0.39\$-0.56\$ per tablet



Lorazepam (Ativan)

#### Common Side Effects

- Confusion
- Drowsiness
- Restlessness
- Irritability
- Tolerance
- Misuse

#### Precautions

- Kidney disease
- Liver disease
- Glaucoma
- Psychosis
- Other substance use



### Barbiturate – Withdrawal Management

Phenobarbital (Luminal)

Route Oral • 15mg, 16.2mg, 30mg, 32.4mg, 60mg, **Formulation** 100mg tabs, 20mg/5ml elixir • Symptom-driven or fixed dose-and-Daily Dose taper protocols Half-Life • 80-120 hours Cost \$15 per tablet; \$50 for 375ml elixir



### Barbiturate – Withdrawal Management

Phenobarbital (Luminal)

## Common Side Effects

- Somnolence
- Sedation
- Restlessness
- Respiratory depression

#### **Precautions**

- Other substance use
- Pregnancy
- Carcinogenic

#### Other

- Narrow therapeutic window
- Considered third-line
- Monitor for toxicity



Carbamazepine (Tegretol)

Route Oral **Formulation** • 100 mg, 200 mg, 300 mg capsules 800mg Day 1 tapering to 200mg Day Daily Dose 4 in divided doses Half-Life • 35-40 hours Cost • 0.20\$-1.21\$ per capsule



Carbamazepine (Tegretol)

#### **Common Side Effects**

- Blurred vision
- Continuous back and forth eye movements
- Diarrhea
- Headaches

#### **Precautions**

- Use of MAO inhibitor
- Psychosis
- History of suicide
- Light sensitivity



Gabapentin (Neurontin)

Route Oral • 100 mg, 300 mg, 400 mg capsules **Formulation** • 600 mg, 800 mg tablets • 600 - 1200 mg divided into 2 to 3 Daily Dose doses daily, tapering over 4 to 7 days Half-Life • 5-7 hours Cost • 0.14\$-4.17\$ per capsule/tab



Gabapentin (Neurontin)

## Common Side Effects

- Ataxia
- Dizziness
- Drowsiness
- Fatigue

#### **Precautions**


- Mental illness
- Seizures
- Pregnancy

#### Other

Physical dependence



Valproic acid (Depakote)





Valproic acid (Depakote)

#### **Common Side Effects**

- Drowsiness
- Diarrhea
- Headache
- Weight changes

#### **Precautions**

- Breastfeeding
- Drowsiness



Acamprosate (Campral)

Route Oral **Formulation** • 333 mg • 2 x 333 mg tab po TID or 3 x 333 mg Daily Dose tab po BID Half-Life • 20-33 hours Cost • \$70-200/month



Acamprosate (Campral)

## Common Side Effects

- Diarrhea
- Nausea
- Depression
- Anxiety

#### **Precautions**

- Dose adjustment for CrCl 30-50ml/min
- Unstable depression or suicidality

#### Other

 Patient selection: Impulsivity issues or protracted withdrawal symptoms



Disulfiram (Antibuse)

Route Oral **Formulation** • 125 mg, 250 mg, 500 mg Daily Dose • 250 mg once daily Half-Life • 60-120 hours Cost • \$54.90/month



Disulfiram (Antibuse)

### Common Side Effects

- Metallic taste
- Dermatitis

#### **Precautions**

- Psychosis
- DM
- Epilepsy
- Hepatic dysfunction
- Thyroid disease
- Renal impairment
- Dermatitis

#### Other

 Patient should be highly motivated and monitored



Naltrexone (ReVia – oral) (Vivitrol – IM)

Route • Oral, IM • 50 mg tab **Formulation** • 380 mg IM • 50mg tab po q day Daily Dose • 380mg IM q month Half-Life • 4-13 hours • Tabs: \$130-300+/month Cost • Single dose vial: \$1500 /month



Naltrexone (ReVia – oral) (Vivitrol – IM)

## Common Side Effects

- Nausea
- Abdominal pain
- Constipation
- Dizziness
- Headache
- Anxiety
- Fatigue

#### **Precautions**

- Liver disease
- Renal impairment
- History of suicide attempts

#### Other

- Patient selection:
  - Family history of alcohol dependency or
  - Concurrent Cognitive Behavioral Therapy (CBT)



Gabapentin (Neurontin)

Route Oral **Formulation** • 200 mg, 300 mg, 400 mg capsules • 600-1200 mg divided into 2 to 3 Daily Dose doses daily, tapering over 4 to 7 days Half-Life • 5-7 hours Cost • \$8-250/month



Gabapentin (Neurontin)

### Common Side Effects

- Ataxia
- Dizziness
- Drowsiness
- Fatigue

#### **Precautions**

- Mental illness
- Seizures
- Pregnancy

#### Other

 Physical dependence and misuse



Topiramate (Topamax)

Route Oral **Formulation** • 25 mg, 50 mg, 100 mg tabs Daily Dose • 300 mg/day po max Half-Life • 19-23 hours Cost • \$9-500/month



Topiramate (Topamax)

## Common Side Effects

- Numbness and tingling
- Change in taste
- Decreased appetite
- Concentration

#### **Precautions**

- Renal impairment with dose adjustment for CrCl
  - < 70ml/min
- Depression
- Glaucoma

#### Other

 Patient selection: Can start non sober



Baclofen (Gablofen, Lioresal)

Route Oral **Formulation** • 5mg, 10mg, 20mg tabs Daily Dose • 60mg-180+ mg daily, in divided doses Half-Life • 2-6 hours Cost • \$60-300/month



Baclofen (Gablofen, Lioresal)

## Common Side Effects

- Drowsiness
- Dizziness
- Weakness
- Nausea
- Confusion
- Urinary retention

#### **Precautions**

- Renal impairment (cleared renally)
- H/o seizure disorder
- Avoid abrupt discontinuation
- Peptic ulcer disease
- Decreases GI motility
- H/o psychosis

#### Other

- Not used as first-line
- Non sober start ok
- Titrate by 10mg q3 days to effective dose